

GOUT

Gout is a disorder that results from the buildup of uric acid in the tissues or a joint. It most often affects the joint of the big toe. Gout attacks are caused by deposits of crystallized uric acid in the joint. Uric acid is present in the blood and eliminated in the urine, but in people who have gout, uric acid accumulates and crystallizes in the joints. Uric acid is the result of the breakdown of purines, chemicals found naturally in our bodies and in food. Some people develop gout because their kidneys have difficulty eliminating normal amounts of uric acid, while others produce too much uric acid.

Gout occurs most commonly in the big toe because uric acid is sensitive to temperature changes. At cooler temperatures, uric acid turns into crystals. Since the toe is the part of the body that is farthest from the heart, it is also the coolest part of the body and, thus, the most likely target of gout. However, gout can affect any joint in the body.

The tendency to accumulate uric acid is often inherited. Other factors that put a person at risk for developing gout include: high blood pressure, diabetes, obesity, surgery, chemotherapy, stress and certain medications and vitamins. For example, the body's ability to remove uric acid can be negatively affected by taking aspirin, some diuretic medications ("water pills") and the vitamin niacin (also called nicotinic acid). While gout is more common in men aged 40 to 60 years, it can occur in younger men as well as in women.

Symptoms

An attack of gout can be miserable, marked by the following symptoms:

- Intense pain that comes on suddenly, often in the middle of the night or upon arising
- Signs of inflammation, such as redness, swelling and warmth over the joint

Diagnosis

To diagnose gout, the foot and ankle surgeon will ask questions about your personal and family medical history, followed by an examination of the affected joint. Laboratory tests and x-rays are sometimes ordered to determine if the inflammation is caused by something other than gout.

Treatment

Initial treatment of an attack of gout typically includes the following:

- Medications. Prescription medications or injections are used to treat the pain, swelling and inflammation.
- Dietary restrictions. Foods and beverages that are high in purines should be avoided since purines are converted in the body to uric acid.

- Fluids. Drink plenty of water and other fluids each day, while also avoiding alcoholic beverages, which cause dehydration.
- Immobilize and elevate the foot. Avoid standing and walking to give your foot a rest. Also, elevate your foot (level with or slightly above the heart) to help reduce swelling.

The symptoms of gout and the inflammatory process usually resolve in three to ten days with treatment. If gout symptoms continue despite the initial treatment, or if repeated attacks occur, see your primary care physician for maintenance treatment that may involve daily medication. In cases of repeated episodes, the underlying problem must be addressed, as the buildup of uric acid over time can cause arthritic damage to the joint.

FLAGSTAFF
BONE & JOINT
ORTHOPAEDIC EXCELLENCE. EXCEPTIONAL CARE.

www.flagstaffboneandjoint.com